

## Corn Yield Response to Nitrogen (0-250 lbs/ac)

Eric Richer, Ohio State University Extension Educator, Fulton County, Ohio  
 Anthony Fulford, Ohio State University Postdoctoral Researcher, Wooster, Ohio  
 Steve Culman, Ohio State University Assistant Professor of Soil Fertility, Wooster, Ohio

### Objective

To determine the effects of total nitrogen rate on corn grain yield.

### Background (all farms)

---

Crop Year: 2015

County: Fulton

Drainage: systematic, 50' laterals or closer

Previous Crop: soybeans

Soil Test: all in maintenance range or higher

Starter: all used a blended N-P analysis

Seeding Rate: 32,000-35,000 seeds/ac

### Methods

This trial evaluates corn yield response to nitrogen. Generally, all participating farmers had similar practices with regard to fertility, herbicide program, traits and soil testing. All plots were replicated at least 4 times (except where noted) in a randomized complete block design. Rates used were 0-250 total units of nitrogen per acre with credits given for starter fertilizer. Plots were field length (approximately 1,200 feet long) except where noted. The farmers' equipment dictated the planting and harvesting width. Each farmer planted/harvested their own trial and applied their own sidedress nitrogen.

### Data

Plot centers were harvested with commercial combines. Yield and moisture data was collected with calibrated yield monitors and shrunk to 15% moisture. Nitrogen use efficiency (NUE) was calculated by dividing total units of nitrogen applied by yield. Normalized Difference Vegetative Index (NDVI) readings were taken with Greenseeker handheld sensors approximately four weeks after nitrogen was applied and corn stalk nitrate tests (CSNT) were taken approximately 1-2 weeks after black layer to evaluate nitrate nitrogen remaining. Weather data was obtained from the closest National Weather Service station or by the farmer.

### Economics

All treatments have economics calculated based on \$3.50/bushel corn marketing price and a cost of \$.46/lb Nitrogen (Source: 2015 OSU Ag Econ Corn Budget). Net return is calculated as gross revenue minus nitrogen cost or as Return less N.



## Results

### OFR Trial: Corn Yield Response to N Rate #1

Location: Delta

Rainfall (May-Sept): 20.4" (producer's records)

Soil Type: Hoytville clay loam/Haskins loam

Tillage: No-till

N Source: NH<sub>3</sub>

	Yield (bu/ac)	Nitrogen Use Efficiency (lb N/bu)	Return less N
0 lbs N/ac	116.0 D	--	\$406/ac
100 lbs N/ac	180.8 C	.55	\$587/ac
150 lbs N/ac	220.3 B	.68	\$702/ac
200 lbs N/ac	250.5 A	.80	\$785/ac
250 lbs N/ac	265.5 A	.94	\$814/ac
LSD (P<.05)	17.9 (CV 4.88)		

### OFR Trial: Corn Yield Response to N Rate #2

Location: Fayette

Rainfall (Apr-Sept): 22.1" (CoCoRaHS)

Soil Type: Fulton silty clay loam

Tillage: No-till

N Source: 28% UAN

	Yield (bu/ac)	NUE (lb N/bu)	Return less N
0 lbs N/ac	73.8 D	--	\$258/ac
100 lbs N/ac	87.0 D	1.2	\$259/ac
150 lbs N/ac	114.4 C	1.3	\$331/ac
200 lbs N/ac	139.7 B	1.4	\$397/ac
250 lbs N/ac	164.4 A	1.5	\$460/ac
LSD (P<.05)	19.7 (CV 9.05)		

### OFR Trial: Corn Yield Response to N Rate #3

Location: Fayette

Rainfall (Apr-Sept): 22.1" (CoCoRaHS)

Soil Type: Fulton silty clay loam

Tillage: No-till

N Source: 28% UAN

	Yield (bu/ac)	NUE (lb N/bu)	Return less N
0 lbs N/ac	73.6 D	--	\$258/ac
100 lbs N/ac	101.1 C	.99	\$308/ac
150 lbs N/ac	119.0 BC	1.3	\$348/ac
200 lbs N/ac	145.2 AB	1.4	\$416/ac
250 lbs N/ac	157.9 A	1.6	\$438/ac
LSD (P<.05)	26.3 (CV 11.7)		



**Discussion:**

Fulton County sustained above average early season rainfall in 2015. Much of this rainfall led to leaching or denitrification of already-applied nitrogen. These trials show a statistically significant difference in yield across the various nitrogen rates. In 2015, the data supports higher rates of nitrogen being applied unless timely late season applications can be made.

Nitrogen Use Efficiency (NUE) was likely affected by early season rainfall. Trials 2 and 3 suggest that yield and economic optimums required NUE greater than 1.0 lbs N per bushel produced. In trial 1, yield and economic optimums were approximately .94 lbs N per bushel produced.

Please refer to 2014 Yield Response to Nitrogen OFR Results in Fulton County ([www.go.osu.edu/fultononfarmresearch](http://www.go.osu.edu/fultononfarmresearch)) and note the importance of multi-year data to determine an economically optimum nitrogen rate you your farm. Generally speaking, 2014 yield response to nitrogen results followed a more average economic optimum nitrogen rate whereas 2015 data supports a higher than average economic optimum nitrogen rate due, in part, to excess rainfall in the early part of the growing season.

**Acknowledgement**

The authors express appreciation to on-farm collaborators Les Seiler and Michael Vorwerk for conducting Corn Yield Response to Nitrogen OFR Plots. Thanks to agronomy intern Troy Grime for assistance with data collection.

